

MYRMELACHISTA (DECAMERA) NODIGERA Mayr

var. FLAVICORNIS Emery

Emery, Zool. Jahrb. Abt. Syst. 9, 1896 p. 638 ♀ fig. F.

Paraguay: San Salvador (J. Bohls), in woody thorns of *Acacia* (probably *cavenia*).

MYRMELACHISTA (DECAMERA) PADEREWSKII Forel

Forel, Verh. zool. bot. Ges. Wien 1908 p. 397 ♀ ♂.

Brazil: São Paulo (E. Lutz); nesting in bamboo.

MYRMELACHISTA (DECAMERA) SCHUMANNI Emery

Emery, Ann. Soc. Ent. France 1890 p. 70 nota ♀; Schumann, Sammlun. gemeinverst. wiss. Vortr. 83, 1889 p. 17; Emery, Biol. Centralbl. 11, 1891, p. 165; Anal. Mus. Nac. Costa Rica 1892 p. 65.

Colombia: (K. Schumann), in cavities of *Duroia hirsuta*.

Var. CORDINCOLA Wheeler & Mann, var. nov.

*Worker.* Length 1.5—3 mm.

Averaging somewhat larger than the typical form of the species and differing in color. Brownish yellow, with the head and gaster brown, the latter usually darker than the former, the bases of the gastric segments paler than their posterior borders which are edged with black. Mandibles and anterior portion of head paler and more yellowish than the posterior portion. Antennæ and legs yellow, the clubs of the former and in some specimens also the middle portions of the femora feebly infuscated. Summit of petiolar node entire, its anterior surface distinctly convex, its posterior surface nearly flat.

Described from numerous specimens taken by Dr. W. M. Mann at Osunto, Bolivia, in the cauline swellings of *Cordia hispidissima*.

MYRMELACHISTA (DECAMERA) ULEI Forel

Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 704 ♀.

Peru: Cerro de Escaler, 1200 m. (E. Ule), in the swellings of the flower stems of a melastomaceous plant.